PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Keep people and pets out of treated areas until sprays have dried. FIRST AID:

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Information, call the National Pesticide Information Center

at 1-800-858-7378

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRODUCT INFORMATION

Treats up to 3,000 Square Feet [Quart] Treats up to 6,000 Square Feet [1/2 Gallon Treats up to 12,000 Square Feet [1 Gallon]

WHERE TO USE:

· Around patios, walkways, driveways and fences.

- In vegetable or flower gardens and around trees and shrubs and other landscaped areas.
- In large areas:

- For lawn renovation or representa-- To prepare garden plots for planting. AMOUNT TO USE: - For lawn renovation or replacement.

For Best Results: 6 fl. oz. (12 Tbs) per gallon of watter. For areas with small or seedling weeds: 3 fl. oz. (6 Tbs) per gallon of water per 300 square

NOTICE: Read the entire label. Use only according to label directions. Before using this product, read "Warranty Disclaimer", "Inherent Risks of Use", and "Limitations of Remedies". If terms are unacceptable, return at once unopened.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

This is an end-use product. The manufacturer does not intend and has not registered it for reformulation.

General Information

This product kills most broadleaf and grass weeds. This product may be used for non-selective weed control around patios, walkways, driveways and fences, in vegetable or flower gardens and around trees and shrubs and other landscaped areas, in large areas for lawn removal prior to replacing turfgrasses, and for weed control in graden plots prior to planting. Applications to plant foliage are rainfast in approximately two hours and afterward will not be affected by rainfall or irrigation.

How This Product Works

The active ingredient in this product is active against a substance found only in plants. It enters the plant through foliage and is translocated throughout the plant, killing both roots and top growth. This product is not active in soil and cannot move through the soil to damage desirable plants. Sprays not absorbed by plants are broken down into natural materials.

What To Expect

Weeds may begin to wilt within a few hours, but complete kill may require 1 to 2 weeks. Small, rapidly growing weeds may be killed with a single application while older, mature weeds may require re-application. If re-application is necessary, wait until new growth appears.

Important Use Precautions

Do not spray during windy conditions or allow the spray to contact desirable plants such as flowers, ground covers or lawn areas, as they will be severely damaged or killed. In situations where weeds and desirable plants are in close proximity, use a piece of cardboard or plastic to shield desirable plants from the spray. If a desirable plant is sprayed by mistake, immediately rinse the plant with water. Before sprays have dried, avoid inadvertent transfer of the spray solution from treated plants to desirable plants or tracking spray from treated areas onto lawns or other desirable vegetation. Do not apply through any type of irrigation system.

Keep people and pets out of treated areas until sprays have dried.

SPRAYER CLEANUP

Rinse sprayer and all sprayer parts three times with water. Apply rinse waters to bare soil or graveled area away from desirable vegetation. After careful cleaning, sprayer may be used for other products.

0155.031610

Printed in U.S.A. 711



United 345

WEED BLITZ

Concentrated Weed Killer For Control of all Types of Broadleaf Weeds and Grasses

For Outdoor Use Only • Easy To Use Kills Weeds, Roots and All • Rainproof in 2 Hours Treats up to 3,000 Square Feet [Quart] Treats up to 6,000 Square Feet [1/2 Gallon] Treats up to 12,000 Square Feet [1 Gallon]

ACTIVE INGREDIENTS:

glyphosate[†]: N-(phosphonomethyl) glycine,

isopropylamine salt	
INERT INGREDIENTS:	
TOTAL INGREDIENTS	
[†] Contains 1.6 pounds	per gallon alvphosate, isopropylamine salt

.o pounds per galion giypnosate, isopropyl

(1.2 pounds per gallon glyphosate acid).

KEEP OUT OF REACH OF CHILDREN. CAUTION

Refer to back panel for Precautionary Statements and Directions for Use, including Storage and Disposal instructions.

EPA REG. NO. 10088-107-9250 EPA EST. 10088-WI-1

Sold By: UNITED LABORATORIES, INC. 320 37th Avenue • St. Charles, IL 60174 1-800-323-2594 • www.united/absinc.com

MIXING INSTRUCTIONS

- Tank Sprayer† (Plastic, aluminum or stainless steel):
- Use 6 fl. oz. (12 Tbs) per gallon of water. Spot treat or spray evenly over 300 sq. ft.

Ortho[®] Dial 'n Spray[®]: • Set dial to 5 fl.oz.

- Use 6 fl. oz. (12 Tbs) per 300 sq. ft. (Do not add water)
- Spray evenly over measured area.
 Tablespoon(Tbs) = 3 teaspoons (tsp)
- fl. oz. = 2 Tbs = 6 tsp

This product can react with certain metal surfaces and should not be applied with galvanized or unlined steel sprayers.

APPLICATION For Best Results

- Apply when weather is warm (above 60°F) and sunny to stimulate systemic movement from foliage to roots.
- To avoid spray drift to desirable vegetation, adjust spray nozzle to produce large droplets and spray when wind is calm.
- Spray foliage until thoroughly wet, but not past the point of runoff.
 Apply when weeds are small and actively growing, before seeds develop.
 Treatment of Vines

To control vines growing on fences, mature trees, or poles, cut the vines back to a height of 3 to 4 feet and spray thoroughly. Avoid spray contact with desirable plants or lawn. For control of vines climbing on trees or shrubs with green bark, remove vines by cutting off at the

base of the plant and spray after the vines have begun to regrow. Shield trees or shrubs with a sheet of cardboard or plastic to prevent contact with spray. Turf Replacement

To kill an existing lawn prior to replacement skip one mowing before spraying and apply at the higher rate to 6 fl. oz. (12 Tbs.) per gallon of water. Control will be better if the grass is actively growing, so water 2 to 3 days before and after application if the soil becomes dry.

Reapply to any green patches that remain after 7 days. Wait an additional 7 days after the last application before raking, tillage prior to reseeding or sodding. Harvesting

If used to control weeds around fruit or nut trees, caneberries, or grapevines, allow 21 days before harvesting.

Replanting

Flowers, shrubs, and trees may be planted 1 day ofter application. **These fruits, vegetables, and herbs may be planted 3 days after application**: Beans, beet greens, broccoli, brussels sprouts, cabbage, carrot, cauliflower, celery, chard (Swiss), chicory, collards, corn, cucumber, eggplant, endive, garlic, gourds, horseradish, Jerusalem artichoke, kale, kohlrabi, leek, lentils, lettuce, melons, mustard greens, okra, onion, rape greens, parsley, parsnips, peas, pepper, potato, pumpkin, radish, rhubarb, rutabaga, spinach, squash, sweet potato, tomatillo, tomato (from

seal), turnip, watercress and yams. Lawn grass and the following caneberries may be planted 7 days after application: Blackberry, blueberry, boysenberry, currant, dewberry, elderberry, gooseberry, huckleberry, loganberry, olallieberry, and raspberry (black and red).

Tomatoes from transplant and unlisted fruits, vegetables and herbs may be planted 30 days after application.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. The use place, term independence backage at once to the same for a full related to particular place place. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies. Warranty Disclaimer Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for

the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO EXPRESS NOR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Lack of performance, or other unintended consequences may occur due to such factors as use of the product contrary to label instructions (including adverse conditions noted on the label, such as unfavorable temperature, soil instructions (including durense containions noted on the table, such as untravorable temperature, soil conditions, etc.), abnormal conditions (such as extreme weather conditions), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

STORAGE AND DISPOSAL Do Not Contaminate Water, Food, or Feed By Storage or Disposal.

Pesticide Storage: Store in a dry place no lower in temperature than 50°F or higher than 120°F. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, sproy mixture or protection and the state of education inductions in important states besincter, sproy mixture, or ninsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Container Disposal: Nonrefillable container. Do not reuse or refil container_Offer for recycling,

if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinstate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times.