DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling This product is not for use on critical and semi-critical medical device surfaces.

CLEANING: Use one United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes to completely remove surface of all aross debris from surfaces to be treated.

DISINFECTING: To disinfect hard, non-porous surfaces, thoroughly wet the surface to be treated. Treated surface must remain visibly wet for disinfection:

- To disinfect against Escherichia coli (E. coli), Esherichia coli O157:H7, Pseudomonas aeruainosa, Salmonella (choleraesuis) enterica and Vancomycin Resistant Enterococcus faecalis (VRE), allow the treated surface to remain wet for 10 minutes.
- To disinfect against Mycobacterium bovis BCG (Tuberculosis), allow the treated surface to remain wet for 10 minutes. • To disinfect against Staphylococcus aureus and Methicillin Resistant Staphylococcus aureus (MRSA), allow the treated surface
- to remain wet for 10 minutes.
- To disinfect against Vancomycin intermediate Resistant Staphylococcus aureus (VISA), allow the treated surface to remain wet for 10 minutes.
- A potable water rinse is required for surfaces in direct contact with food.

For plastic and painted surfaces, spot test on an inconspicuous area before use.

For heavily soiled surfaces, clean before following disinfecting instructions.

VIRUCIDAL ACTIVITY: United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes are an effective virucide on hard, non-porous surfaces against: • Human Immunodeficiency Virus Type 1 (HIV-1) when the treated surface is allowed to remain wet for 2 minutes. • Norovirus (Norwalk Virus) and Feline calicivirus when the treated surface is allowed to remain wet for 10 minutes. Influenza A (H1N1) Virus, (ATCC VR 1469, Strain A/PR/8/34) when the treated surface is allowed to remain wet for 10 minutes. Use additional wipes if needed to assure that the treated surface remains continuously wet for the prescribed time.

For Ultrasound Transducers and Probes:

Use only on external surfaces of ultrasound transducers and probes. With a clean washcloth, clean ael from external surface of the ultrasound transducer or probe to be disinfected. Then thoroughly wet probe with wipe. Treated surface must remain wet for 10 minutes. Use additional wipes if needed to assure continuous 10 minute wet contact time. Let air dry

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/ BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AÍDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/ OBJECTS SOILED WITH BLOOD / BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be throoughly cleaned from surfaces and objects before application of this product. United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes can be used for this purpose.

CONTACT TIME: Thoroughly wet surface with a new wipe. Allow surface to air dry. Although efficacy at a 2 minute contact time has been shown to be adequate against HIV-1, use the contact time for the disinfection of other organisms listed on this label. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal

Not for personal cleansing. This is **not** a baby wipe! Do not flush in toilet.

TO OPEN:

1. Remove lid from canister

2. Lift flap in center of lid.

3. Pull up corner of center wipe, twist to a point and thread through dispenser hole in lid. DO NOT PUSH FINGER THROUGH OPENING

4. Replace lid. Pull out exposed wipe and snap off. The next wipe pops up automatically. Dispose of used wipes in trash receptacle. Snap center flap down when finished to retain moisture.

5. Do not flush



United 368

HARD-D-WIPES Hard Surface Disinfectant Wipes

FAST EFFECTIVE ANTIBACTERIAL FORMULA CLEANS AND DISINFECTS BLEACH-FREE • ALCOHOL-FREE • *VIRUCIDAL TUBERCULOCIDAL · BACTERICIDAL **PSEUDOMONICIDAL**

Kills Pandemic 2009 HINI Influenza A virus (formerly called Swine Flu)

ACTIVE INGREDIENTS	
Alkyl (60% C 14, 30% C16 , 5% C12 , 5% C18)	
dimethyl benzyl ammonium chlorides	0.105%
Alkyl (68% C 12, 32% C14)	
dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS	99.790 %
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements.

EPA REG. NO. 1839-174-9250 EPA EST. NO. 10088-WI-2 NET CONTENTS: 2 lbs. 10.5 ozs. 180 Pre-measured Wipes 6" x 7" (15.24cm x 17.78cm)

Sold By: UNITED LABORATORIES, INC. 320 37th Avenue • St. Charles, IL 60174 • 1-800-323-2594 • www.unitedlabsinc.com

United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes are intended for use in: Hospitals and healthcare settings: ambulances, anesthesia, CAT labs, CCU, Central Supply, Clinics, Dental Offices, Dialysis Clinics, Doctor's office(s), Emergency Rooms, E.R., Home health care, Hospices, ICU, Laboratory, Laundry rooms, Neonatal Intensive Care Units (NICU), Newborn nursery, Nursing homes, Operating rooms, Pediatrics, Pediatric Intensive Care Units (PICU), Physical therapy, Physician's office(s), Opthalmic offices, Radiology, Recovery Room, Rehabilitation, Respiratory therapy, Surgery, Transport vehicles, X-Ray; Critical Care Areas: CCU, Emergency Rooms, E.R., I.C.U., Neonatal Intensive Care Units (NICU), Operating Rooms, Pediatric Intensive care Units (PICU), Surgery, Opthalmic offices; Veterinary clinics, animal hospitals, animal life science laboratories, kennels, pet animal kennels, breeding establishments, grooming establishments, pet animal quarters, zoos, pet shops, tack shops, animal holding areas, veterinary waiting rooms, veterinary examination rooms, veterinary X-ray rooms, veterinary operating rooms, animal transportation vehicles, beauty salons, barber shops, correctional facilities, funeral homes, mortuaries, schools, health clubs, gyms, athletic facilities, airports, transportation terminals, Grocery stores, Shopping Centers, Delicatessens, Catering facilities, Banquet halls, Nursery, Daycare centers. United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes will clean and disinfect hard, non-porous, surfaces of: bed railings, cabinets, carts, chairs, counters, exam tables, gurneys, isolettes, IV poles, stethoscopes, stretchers, tables, telephones, toilet seats, toilets, bathtubs, bathroom fixtures, shower stalls, sinks, tiles, vanity tops, hospital beds, traction devices, MRI, CAT, scales, paddles, wheelchairs, ultrasound transducers and probes, grocery cart handles, grocery carts, shopping carts, grocery cart child seats, changing tables, cribs, diaper pails, trashcans, garbage cans, cabinets, chairs, computers, countertops, desks, desktops, doorknobs, fax machines, headsets, keyboards, railings, seats, sinks, tables, tabletops, telephones, gym equipment, exercise machines, wrestling mats, and

gymnastic equipment. Use United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes to clean and disinfect non-porous personal protective safety equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, silicone rubber or PVC hearing protectors and vinyl covered ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Caution: Cleaning at the 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

United 368 HARD-D-WIPES Hard Surface Disinfectant Wipes are formulated to disinfect hard, non-porous, inanimate environmental surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, nonpolycarbonate plastic surfaces, bathrooms, shower stalls, bathtubs, and cabinets.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE: Store in an area inaccessible to children. CONTAINER STORAGE/DISPOSAL: Wipes: Dispose of wipe in trash after use. Do not flush. Canister: Tightly close lid between uses to retain moisture. Nonrefillable container. Do not reuse or refill this canister. Offer empty canister for recycling if available or discard in trash.

PRECAUTIONARY STATEMENTS Hazards To Humans And Domestic Animals

CAUTION:

eating, drinking, chewing gum or using tobacco.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. You may contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

1616.090110

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before

FIRST AID

Printed in U.S.A.

1213